


The CDC says you should get a flu vaccine as long as flu viruses are circulating. And since flu season can last as late as May, you still have time.

Help protect yourself and eligible family members ages 2-49 against the flu with FluMist® Quadrivalent



FLU SEASON IS HERE, BUT... IT'S NOT TOO LATE TO VACCINATE

Did you know that the flu typically peaks in February?



Talk to your pharmacist about FluMist Quadrivalent, the only needle-free, nasal-spray flu vaccine that starts working in the nose, where the flu typically starts

Select Safety and Eligibility Information

FluMist Quadrivalent is a vaccine that is sprayed into the nose to help protect against influenza. It can be used in eligible children, adolescents, and adults ages 2 through 49. FluMist Quadrivalent may not prevent influenza in everyone who gets vaccinated. FluMist Quadrivalent is for intranasal administration only.

You should not get FluMist Quadrivalent if you have a severe allergy to eggs or to any inactive ingredient in the vaccine; have ever had a life-threatening reaction to influenza vaccinations; or are 2 through 17 years old and take aspirin or medicines containing aspirin – children or adolescents should not be given aspirin for 4 weeks after getting FluMist Quadrivalent unless your healthcare provider tells you otherwise.

Please see additional Important Safety and Eligibility Information on reverse side.




FluMist® Quadrivalent
Influenza Vaccine Live, Intranasal

FluMistQuadrivalent.com

Important Safety and Eligibility Information

What is FluMist® Quadrivalent (Influenza Vaccine Live, Intranasal)?

FluMist Quadrivalent is a vaccine that is sprayed into the nose to help protect against influenza. It can be used in children, adolescents, and adults ages 2 through 49. FluMist Quadrivalent is similar to MedImmune's trivalent influenza vaccine, except FluMist Quadrivalent provides protection against an additional influenza strain. FluMist Quadrivalent may not prevent influenza in everyone who gets vaccinated.

Who should not get FluMist Quadrivalent?

You should not get FluMist Quadrivalent if you have a severe allergy to eggs or to any inactive ingredient in the vaccine; have ever had a life-threatening reaction to influenza vaccinations; or are 2 through 17 years old and take aspirin or medicines containing aspirin – children or adolescents should not be given aspirin for 4 weeks after getting FluMist Quadrivalent unless your healthcare provider tells you otherwise.

Children under 2 years old have an increased risk of wheezing (difficulty with breathing) after getting FluMist Quadrivalent.

Who may not be able to get FluMist Quadrivalent?

Tell your healthcare provider if you or your child are currently wheezing; have a history of wheezing if under 5 years old; have had Guillain-Barré syndrome; have a weakened immune system or live with someone who has a severely weakened immune system; have problems with your heart, kidneys, or lungs; have diabetes; are pregnant or nursing; or are taking Tamiflu®, Relenza®, amantadine, or rimantadine.

Your healthcare provider will decide if FluMist Quadrivalent is right for you or your child.

What are the most common side effects of FluMist Quadrivalent?

The most common side effects are runny or stuffy nose, sore throat, and fever over 100°F.

Please see accompanying complete Product Information, including Patient Information.

Tamiflu and Relenza are registered trademarks of their respective owners.

